

New Products - Priligy

Priligy (**dapoxetine hydrochloride**) is a SSRI. Priligy is indicated for the treatment of premature ejaculation (PE) in men 18 to 64 years of age, who have all of the following: an intravaginal ejaculatory latency time (IELT) of less than two minutes; a persistent or recurrent ejaculation with minimal sexual stimulation before, on, or shortly after penetration and before the patient wishes; a marked personal distress or interpersonal difficulty as a consequence of PE; and a poor control over ejaculation. Priligy is contraindicated in all of the following conditions: significant pathological cardiac conditions (eg NYHA class II-IV heart failure), conduction abnormality (2nd, 3rd degree A-V block or sick sinus syndrome) not treated with permanent pacemaker, significant ischaemic heart, valvular disease; concomitant MAOIs, thioridazine, SSRIs, SNRIs, TCAs, serotonergic drugs/ herbs (eg L-tryptophan, triptans, tramadol, linezolid, lithium, St John's wort) (including less than 14 days before starting, less than 7 days after ceasing dapoxetine); concomitant potent CYP3A4 inhibitors eg ketoconazole, itraconazole, atazanavir, nelfinavir, ritonavir, saquinavir, telithromycin, nefazodone; and in moderate to severe hepatic impairment. Priligy is available as a 30 mg tablet in packs of 3's and 6's.

This list is a summary of only some of the changes that have occurred over the last month. Before prescribing, always refer to the full product information.